arXiv:0704.1549v1 [math.OA] 12 Apr 2007

SATURATED ACTIONS BY FINITE DIMENSIONAL HOPF $*$-ALGEBRAS ON C^*-ALGEBRAS

JA A JEONG[†] AND GI HYUN PARK[‡]

Abstract. If a finite group action α on a unital C^*-algebra M is saturated, the canonical conditional expectation $E : M \to M^\alpha$ onto the fixed point algebra is known to be of index finite type with $Index(E) = |G|$ in the sense of Watatani. More generally if a finite dimensional Hopf $*$-algebra A acts on M and the action is saturated, the same is true with $Index(E) = \dim(A)$. In this paper we prove that the converse is true. Especially in case M is a commutative C^*-algebra $C(X)$ and α is a finite group action, we give an equivalent condition in order that the expectation $E : C(X) \to C(X)^\alpha$ is of index finite type, from which we obtain that α is saturated if and only if G acts freely on X. Actions by compact groups are also considered to show that the gauge action γ on a graph C^*-algebra $C^*(E)$ associated with a locally finite directed graph E is saturated.

1. Introduction

It is known [17] that if α is an action by a compact group G on a C^*-algebra M, the fixed point algebra M^α is isomorphic to a hereditary subalgebra $e(M \times_\alpha G)e$ of the crossed product $M \times_\alpha G$ for a projection e in the multiplier algebra of $M \times_\alpha G$. If $e(M \times_\alpha G)e$ is full in $M \times_\alpha G$ (that is, $e(M \times_\alpha G)e$ generates $M \times_\alpha G$ as a closed two-sided ideal), the action α is said to be saturated (the notion of saturated action was introduced by Rieffel [14, Chap.7]). Every action α with a simple crossed product $M \times_\alpha G$ is obviously saturated.

On the other hand, an action of a finite dimensional Hopf $*$-algebra A on a unital C^*-algebra M is considered in [18] and it is shown that if the action is saturated, the canonical conditional expectation $E : M \to M^A$ onto the fixed point algebra M^A is of index finite type in the sense of Watatani [19] and $Index(E) = (\dim A)1$. The main purpose of the present paper is to prove that the converse is also true. We see from our result that for an action α by a finite group G, α is saturated if and only if the canonical expectation $E : M \to M^\alpha$ is of index-finite type with index $Index(E) = |G|$.

Besides, we consider actions by compact groups to study the saturation property of a gauge action γ on a C^*-algebra $C^*(E)$ associated with a locally finite directed graph E with no sinks or sources. This paper is organized as follows.

In section 2, we review the C^*-basic construction from [19] and finite dimensional Hopf $*$-algebras from [18] setting up our notations. Then we prove in section 3 that if A is a finite dimensional Hopf $*$-algebra acting on a unital C^*-algebra M such that $E : M \to M^A$ is of index finite type with $Index(E) = (dimA)1$, then the action is saturated (Theorem 3.3).

In section 4, we deal with the crossed product $M \times_\alpha G$ by a finite group in detail and give other equivalent conditions in order that α be saturated. From the

---

Research supported by KRF-ABRL-R14(2003-2008)[†] and Hanshin University Research Grant[‡].

conditions one easily see that an action with the Rokhlin property [7] is always saturated. Also we shall show that if M has the cancellation, an action with the tracial Rokhlin property [12] on M is saturated.

Note that even for an action α by the finite group $\mathbb{Z}_2$, the expectation $E : M \to M^\alpha$ may not be of index finite type in general [19, Example 2.8.4]. For a commutative C^*-algebra $C(X)$ and a finite group action α, we give a necessary and sufficient condition that $E : C(X) \to C(X)^\alpha$ is of index finite type (Theorem 4.10) and provide a formula for $Index(E)$. Then as a corollary we obtain that α is saturated if and only if G acts freely on X.

In section 5, we consider a compact group action α and investigate the ideal $\mathcal{J}_\alpha$ of $M \times_\alpha G$ generated by the hereditary subalgebra $e(M \times_\alpha G)e$. Then we apply the result on $\mathcal{J}_\alpha$ to the gauge action on a graph C^*-algebra in section 6. As a generalization of the Cuntz-Krieger algebras [5], the class of graph C^*-algebras $C^*(E)$ associated with directed graphs E has been studied in various directions by considerably many authors (for example see the bibliography in the book [15] by Raeburn). In [9], Kumjian and Pask show among others that if γ is the gauge action on $C^*(E)$, then $C^*(E)^\gamma$ is stably isomorphic to the crossed product $C^*(E) \times_\gamma \mathbb{T}$, which was done by hiring the notions of skew product of graphs and groupoid C^*-algebras. In Theorem 6.3 we shall directly show that the gauge action is actually saturated (this implies that $C^*(E)^\gamma$ and $C^*(E) \times_\gamma \mathbb{T}$ are stably isomorphic).

2. PRELIMINARIES

Watatani's index theory for C^*-algebras. In [19], Watatani developed the index theory for C^*-algebras, and here we briefly review the basic construction $C^*(B, e_A)$. Let B be a C^*-algebra and A its C^*-subalgebra containing the unit of B. Let $E : B \to A$ be a faithful conditional expectation. If there exist finitely many elements $\{v_i\}_{i=1}^n$ in B satisfying the following

$$b = \sum_i E(bv_i)v_i^* = \sum_i v_i E(v_i^* b), \quad \text{for every } b \in B,$$

E is said to be of *index-finite type* and $\{(v_i, v_i^*)\}_{i=1}^n$ is called a *quasi-basis* for E. The positive element $\sum_i v_i v_i^*$ is then the *index* of E, $Index(E)$, which is known to be an element in the center of B and does not depend on the choice of quasi-bases for E ([19, Proposition 1.2.8]). Let $\mathcal{B}$ be the completion of the pre-Hilbert module $\mathcal{B}_0 = \{\eta(b) \mid b \in B\}$ over A with an A-valued inner product

$$\langle \eta(x), \eta(y) \rangle = E(x^* y), \ \eta(x), \eta(y) \in \mathcal{B}_0.$$

Let $\mathcal{L}_A(\mathcal{B})$ be the C^*-algebra of all (right) A-module homomorphisms on $\mathcal{B}$ with adjoints. For $T \in \mathcal{L}_A(\mathcal{B})$, the norm $\|T\| = \sup\{\|Tx\| : \|x\| = 1\}$ is always bounded. Each $b \in B$ is regarded as an operator L_b in $\mathcal{L}_A(\mathcal{B})$ defined by $L_b(\eta(x)) = \eta(bx)$ for $\eta(x) \in \mathcal{B}_0$. By $e_A : \mathcal{B} \to \mathcal{B}$ we denote the projection in $\mathcal{L}_A(\mathcal{B})$ such that $e_A(\eta(x)) = \eta(E(x))$, $\eta(x) \in \mathcal{B}_0$. Then the *$C^*$-basic construction* $C^*(B, e_A)$ is the C^*-subalgebra of $\mathcal{L}_A(\mathcal{B})$ in which the linear span of elements $L_b e_A L_{b'}$ $(b, b' \in B)$ is dense.

Finite dimensional Hopf $*$-algebras. As in [18], a finite dimensional Hopf $*$-algebra is a finite matrix pseudogroup of [20]. We review from [18] the definition and some basic properties of a finite dimensional Hopf $*$-algebra which we need in the following section.

Definition 2.1. ([18, Proposition 2.1]) A finite dimensional C^*-algebra is called a *finite dimensional Hopf $*$-algebra* if there exist three linear maps,

$$\Delta : A \to A \otimes A, \quad \epsilon : A \to \mathbb{C}, \quad \text{and } S : A \to A$$

which satisfy the following properties

(i) Δ(comultiplication) and ϵ(counit) are $*$-homomorphisms, and S(antipode) is a $*$-preserving antimultiplicative involution,
(ii) $\Delta(1) = 1 \otimes 1$, $\epsilon(1) = 1$, $S(1) = 1$,
(iii) $(\Delta \otimes id)\Delta = (id \otimes \Delta)\Delta$,
(iv) $(\epsilon \otimes id)\Delta = \Delta(\epsilon \otimes id)$,
(v) $m(S \otimes id)(\Delta(a)) = \epsilon(a)1 = m(id \otimes S)(\Delta(a))$ for $a \in A$, where $m : A \otimes A \to A$ is the multiplication.

Proposition 2.2. ([18], [20]) *Let A be a finite dimensional Hopf $*$-algebra. Then the following properties hold.*

(i) *For $a \in A$, with the notation $\Delta(a) = \sum_i a_i^L \otimes a_i^R$, we have*

$$\begin{aligned} \sum_i \epsilon(a_i^L) a_i^R &= a = \sum_i \epsilon(a_i^R) a_i^L, \\ \sum_i a_i^L S(a_i^R) &= \epsilon(a)1 = \sum_i S(a_i^L) a_i^R, \\ \sum_i a_i^R S(a_i^L) &= \epsilon(a)1 = \sum_i S(a_i^R) a_i^L. \end{aligned}$$

(ii) *There is a unique normalized trace (called the Haar trace) τ on A such that*

$$\sum_i \tau(a_i^L) a_i^R = \tau(a)1 = \sum_i \tau(a_i^R) a_i^L, \ a \in A.$$

(iii) *There exists a minimal central projection $e \in A$ (called the distinguished projection) such that $ae = \epsilon(a)e$, $a \in A$. We have*

$$\epsilon(a) = 1, \ S(e) = e, \ \textit{and} \ \ \tau(e) = (\dim A)^{-1}.$$

3. Actions by finite dimensional Hopf $*$-algebras

Throughout this section A will be a finite dimensional Hopf $*$-algebra. An *action* of A on a unital C^*-algebra M is a bilinear map $\cdot : A \times M \to M$ such that for $a, b \in A$, $x, y \in M$,

$$\begin{aligned} 1 \cdot x &= x, \\ a \cdot 1 &= \epsilon(a)1, \\ ab \cdot x &= a \cdot (b \cdot x), \\ a \cdot xy &= \sum_i (a_i^L \cdot x)(a_i^R \cdot y), \\ (a \cdot x)^* &= S(a^*) \cdot x^*. \end{aligned}$$

Then the *crossed product* $M \rtimes A$ is the algebraic tensor product $M \otimes A$ as a vector space with the following multiplication and $*$-operation:

$$(x \otimes a)(y \otimes b) := \sum_j x(a_i^L \cdot y) \otimes a_i^R b,$$

$$(x \otimes a)^* := \sum_i (a_i^L)^* \cdot x^* \otimes (a_i^R)^*.$$

Identifying $a \in A$ with $1 \otimes a$ and $x \in M$ with $x \otimes 1$, we see [18] that

$$M \rtimes A = span\{xa \mid x \in M,\, a \in A\}.$$

For the definition of saturated action of A on M, refer to section 4 of [18].

Proposition 3.1. ([18]) *Let $M^A = \{x \in M \mid a \cdot x = \epsilon(a)x, \text{ for all } a \in A\}$ be the fixed point algebra for the action of A on a unital C^*-algebra M.*

(i) *The action is saturated if and only if*

$$M \rtimes A = span\{xey \mid x, y \in M\},$$

where $e \in A$ is the distinguished projection.

(ii) *The map $E : M \to M^A$, $E(x) = e \cdot x$, is a faithful conditional expectation onto the fixed point algebra such that*

$$E((a \cdot x)y) = E(x(S(a) \cdot y)), \quad a \in A, \; x, y \in M.$$

(iii) *The linear map $F : M \rtimes A \to M$, $F(xa) = \tau(a)x$, is a faithful conditional expectation onto M.*

Recall that $\mathcal{M}_0 := M$ is an M^A-valued inner product module by

$$\langle \eta(x), \eta(y) \rangle_{M^A} = E(x^*y)$$

(here we use the convention in [19] for the inner product as in section 2). Since every norm bounded M^A-module map on $\mathcal{M}_0$ extends uniquely to the Hilbert M^A-module $\mathcal{M}$, we may identify the $*$-algebra $End(\mathcal{M}_0)$ (in [18]) of norm bounded right M^A-module endomorphisms of $\mathcal{M}_0$ having an adjoint with the C^*-algebra $\mathcal{L}_{M^A}(\mathcal{M})$ explained in section 2.

Remark 3.2. ([19, Proposition 1.3.3]) If $E : M \to M^A$ is of index-finite type, then

$$C^*(M, e_{M^A}) = span\{L_x e_{M^A} L_y \mid x, y \in M\} = \mathcal{L}_{M^A}(\mathcal{M}).$$

In fact, we see from the proof of [19, Proposition 2.1.5] that $C^*(M, e_{M^A})$ contains the unit of $\mathcal{L}_{M^A}(\mathcal{M})$. Thus the ideal $span\{L_x e_{M^A} L_y \mid x, y \in M\}$ which is dense in $C^*(M, e_{M^A})$ must contain the unit of $\mathcal{L}_{M^A}(\mathcal{M})$.

Theorem 3.3. *Let A be a finite dimensional Hopf $*$-algebra acting on a unital C^*-algebra M. Then the following are equivalent:*

(i) *The action is saturated.*

(ii) *$E : M \to M^A$ is of index finite type with $Index(E) = (dimA)1$.*

Proof. (i)$\Longrightarrow$ (ii) is shown in [18, Proposition 4.5].
(ii)$\Longrightarrow$ (i). By Remark 3.2, $C^*(M, e_{M^A}) = span\{L_x e_{M^A} L_y \mid x, y \in M\}$. Consider a map $\varphi : C^*(M, e_{M^A}) \to M \rtimes A$ given by

$$\varphi(\sum_i L_{x_i} e_{M^A} L_{y_i}) = \sum_i x_i e y_i.$$

To see that φ is well defined, let $\sum_i L_{x_i} e_{M^A} L_{y_i} = 0$. Then for each $z \in M$,

$$(\sum_i L_{x_i} e_{M^A} L_{y_i})(\eta(z)) = \sum_i \eta(x_i E(y_i z)) = \eta(\sum_i x_i(e \cdot (y_i z))) = 0,$$

hence by the injectivity of η ([19, 2.1]), $\sum_i x_i(e\cdot(y_i z)) = 0$ in M. Since $(a\cdot x)e = axe$ for $a \in A$, $a \in M$ (see (7) of [18]), we thus have

$$\sum_i x_i(e \cdot (y_i z))e = \sum_i (x_i e y_i) z e = 0$$

in $M \rtimes A$ for every $z \in M$, which then implies that

$$(\sum_i x_i e y_i)(z e z') = 0, \;\; z, z' \in M.$$

Particulary, $(\sum_i x_i e y_i)(\sum_i x_i e y_i)^* = 0$, so that $\sum_i x_i e y_i = 0$ (in $M \rtimes A$). Thus φ is well defined. It is tedious to show that φ is a $*$-homomorphism such that the range $\varphi(C^*(M, e_{M^A})) = MeM$ is an ideal of $M \rtimes A$; if x, y, and $z \in M$ and $a \in A$, then

$$(za)(xey) = (z(a \cdot x))ey \in MeM.$$

Hence it suffices to show that $\varphi(1) = 1$. If $\{(u_i, u_i^*)\}_{i=1}^n$ is a quasi-basis for E, then

$$\sum_i L_{u_i} e_{M^A} L_{u_i^*}(\eta(z)) = \sum_i \eta(u_i E(u_i^* z)) = \eta(z), \quad z \in M,$$

which means that $\sum_i L_{u_i} e_{M^A} L_{u_i^*} = 1 \in \mathcal{L}_{M^A}(\mathcal{M})$. Therefore by Proposition 2.2(iii) and Proposition 3.1(iii)

$$F(\varphi(1)) = F(\sum_{i=1}^n u_i e u_i^*) = \sum_i \tau(e) u_i u_i^* = \frac{1}{\dim A} \sum_i u_i u_i^* = 1.$$

Since φ is a $*$-homomorphism, $\varphi(1)$ is a projection in $M \rtimes A$ such that $F(1 - \varphi(1)) = 0$. But F is faithful, and $\varphi(1) = 1$ follows. □

4. ACTIONS BY FINITE GROUPS

Throughout this section G will denote a finite group. As is well known the group C^*-algebra $C^*(G)$ generated by the unitaries $\{\lambda_g \mid g \in G\}$ is a finite dimensional Hopf $*$-algebra with

$$\Delta(\lambda_g) = \lambda_g \otimes \lambda_g, \;\; \epsilon(\lambda_g) = 1, \;\; S(\lambda_g) = \lambda_{g^{-1}} \;\text{ for } \lambda_g \in C^*(G).$$

The Haar trace τ is given by $\tau(\lambda_g) = \delta_{\iota g}$, where ι is the identity of G, and the distinguished projection is $e = \frac{1}{|G|}\sum_g \lambda_g$.

Let α be an action of G on a unital C^*-algebra M. Then it is easy to see that

$$\lambda_g \cdot x := \alpha_g(x) \;\text{ for } g \in G, \; x \in M,$$

defines an action of $C^*(G)$ on M. Furthermore $M \rtimes C^*(G)$ is nothing but the usual crossed product $M \times_\alpha G = span\{x\lambda_g \mid x \in M,\ g \in G\}$, and the expectations $E : M \to M^\alpha (= M^{C^*(G)})$, $F : M \times_\alpha G \to M$ of Proposition 3.1 are given by

$$E(x) = \frac{1}{|G|}\sum_g \alpha_g(x) \ \text{ and } \ F(\sum_g x_g\lambda_g) = x_\iota \ (x, x_g \in M,\ g \in G). \tag{1}$$

Note that for each $\sum_h x_h\lambda_h \in M \times_\alpha G$ and $g \in G$,

$$\|x_g\| = \|F((\sum_h x_h\lambda_h)\lambda_{g^{-1}})\| \le \|(\sum_h x_h\lambda_h)\lambda_{g^{-1}}\| = \|\sum_h x_h\lambda_h\|. \tag{2}$$

If $\mathcal{J}_\alpha$ denotes the closed ideal of $M \times_\alpha G$ generated by the distinguished projection e, then Proposition 3.1(i) says that α is saturated if and only if $\mathcal{J}_\alpha = M \times_\alpha G$. We will see in Proposition 5.4 that

$$\mathcal{J}_\alpha \ = \ span\{\sum_g x\alpha_g(y)\lambda_g \mid x, y \in M\} \ = \ span\{\sum_g x\alpha_g(x^*)\lambda_g \mid x \in M\}. \tag{3}$$

The $*$-homomorphism $\varphi : C^*(M, e_{M^\alpha}) \to M \times_\alpha G$ we discussed in the proof of Theorem 3.3 can be rewritten as follows.

$$\varphi(L_x e_{M^\alpha} L_y) = \frac{1}{|G|}\sum_g x\alpha_g(y)\lambda_g,\ x, y \in M \tag{4}$$

because $\varphi(L_x e_{M^\alpha} L_y) = xey$ and $e = \frac{1}{|G|}\sum_g \lambda_g$. If $\{(u_i, u_i^*)\}$ is a quasi-basis for E, we see from $\sum_i L_{u_i} e_{M^\alpha} L_{u_i^*} = 1$ and (4) that

$$\varphi(1) = \sum_i u_i e u_i^* = \frac{1}{|G|}\sum_g (\sum_i u_i\alpha_g(u_i^*))\lambda_g \tag{5}$$

is a projection in $M \times_\alpha G$. Recall that $\varphi(1) = 1$ holds if α is saturated.

Theorem 4.1. *Let M be a unital C^*-algebra and α be an action of a finite group G on M. Then the following are equivalent:*

- (i) *α is saturated, that is, $\mathcal{J}_\alpha = M \times_\alpha G$.*
- (ii) *$E : M \to M^\alpha$ is of index finite type with $Index(E) = |G|$.*
- (iii) *$E : M \to M^\alpha$ is of index finite type with $Index(E) = |G|$ and*

$$\sum_i u_i\alpha_g(u_i^*) = 0,\ g \neq \iota \tag{6}$$

  for a quasi-basis $\{(u_i, u_i^*)\}$ for E.
- (iv) *There exist $\{b_g^j \in M \mid g \in G,\ 1 \le j \le m\}$ for some $m \ge 1$ such that*
  - (a) *$\alpha_g(b_h^j) = b_{gh}^j$, for $j = 1, \ldots, m$ and $g, h \in G$.*
  - (b) *$\sum_j b_g^j (b_h^j)^* = \delta_{gh}$.*
- (v) *For every $\varepsilon > 0$, there exist $\{b_g^j \in M \mid g \in G,\ 1 \le j \le m\}$ for some $m \ge 1$ such that*
  - (a) *$\sum_j \|\alpha_g(b_h^j) - b_{gh}^j\| < \varepsilon$,*
  - (b) *$\|\sum_j b_g^j (b_h^j)^* - \delta_{gh}\| < \varepsilon$.*

Proof. (i) $\Longleftrightarrow$ (ii) follows from Theorem 3.3.
(i) $\Longrightarrow$ (iii). If $\{(u_i, u_i^*)\}$ is a quasi-basis for E, we have from (5) that $\sum_i u_i\alpha_g(u_i^*) = 0$ for $g \neq \iota$ since $\varphi(1) = 1$.

(iii) $\Longrightarrow$ (ii). Obvious.

(i) $\Longrightarrow$ (iv). Suppose $\mathcal{J}_\alpha = M \times_\alpha G$. By (3) there exist $m \in \mathbb{N}$ and $b_j \in M$, $1 \le j \le m$, such that

$$\sum_g \big(\sum_j b_j \alpha_g(b_j^*)\big)\lambda_g = 1.$$

Thus

$$\sum_j b_j b_j^* = 1 \ \text{ and } \sum_j b_j \alpha_g(b_j^*) = 0 \ \text{ for } g \neq \iota. \tag{7}$$

Set $b_g^j := \alpha_g(b_j)$. Then

$$\begin{aligned} \alpha_g(b_h^j) &= \alpha_g(\alpha_h(b_j)) = \alpha_{gh}(b_j) = b_{gh}^j, \\ \sum_j b_g^j (b_h^j)^* &= \sum_j \alpha_g(b_j)\alpha_h(b_j^*) \\ &= \alpha_g\big(\sum_j b_j \alpha_{g^{-1}h}(b_j^*)\big) \\ &= \delta_{gh} \text{ by (7)}. \end{aligned}$$

(iv) $\Longrightarrow$ (v). Obvious.

(v) $\Longrightarrow$ (i). Let $\varepsilon > 0$ and let $\{b_g^j \in M \mid g \in G,\ 1 \le j \le m\}$ satisfy (a) and (b) of (v). Note that (b) implies $\|b_g^j\| < 1 + \varepsilon$ for $g \in G,\ 1 \le j \le m$. Indeed from $\big\| \|\sum_j b_g^j (b_g^j)^*\| - 1 \big\| \le \|\sum_j b_g^j (b_g^j)^* - 1\| < \varepsilon$, we have $\|b_g^j\|^2 \le \|\sum_j b_g^j (b_g^j)^*\| < 1 + \varepsilon$. Then

$$\begin{aligned} &\| \sum_{h,j} (\sum_g b_h^j \alpha_g((b_h^j)^*)\lambda_g) - |G| \, \| \\ &= \| \sum_g (\sum_{h,j} b_h^j \alpha_g((b_h^j)^*)\lambda_g - |G| \, \| \\ &\le \| \sum_{h,j} b_h^j (b_h^j)^* - |G| \, \| + \| \sum_{g \neq \iota} (\sum_{h,j} b_h^j \alpha_g((b_h^j)^*)\lambda_g) \| \\ &\le \ \sum_h \| \sum_j b_h^j (b_h^j)^* - 1 \, \| + \sum_{g \neq \iota} \sum_h \| \sum_j b_h^j \alpha_g((b_h^j)^*) \| \\ &< \varepsilon |G| + \sum_{g \neq \iota} \sum_h \| \sum_j b_h^j \big(\alpha_g((b_h^j)^*) - (b_{gh}^j)^*\big) \| + \sum_{g \neq \iota} \sum_h \| \sum_j b_h^j (b_{gh}^j)^* \| \\ &< \varepsilon \Big(|G| + |G|^2 \max_{g,j} \|b_g^j\| + |G|^2 \Big) \\ &< \varepsilon \Big(|G| + |G|^2 (1 + \varepsilon) + |G|^2 \Big). \end{aligned}$$

Since $\sum_{h,j} (\sum_g b_h^j \alpha_g((b_h^j)^*)\lambda_g) \in \mathcal{J}_\alpha$ and ε can be chosen to be arbitrarily small, we conclude that $\mathcal{J}_\alpha = M \times_\alpha G$. $\square$

Example 4.2. Let $w = \begin{pmatrix} z_1 & 0 \\ 0 & z_2 \end{pmatrix}$ be a unitary with $w^n = 1$ and define an automorphism α on $M_2(\mathbb{C}))$ by $\alpha(a) = waw^*$, $a \in M_2(\mathbb{C})$. We will show that α is saturated if and only if $z_2 = -z_1$. For this, recall from (3) that α is saturated if and only if there exist $x_j \in M_2(\mathbb{C})$, $1 \le i \le m$, satisfying

$$\sum_{k=0}^{n-1} \sum_{j=1}^{m} x_j \alpha^k(x_j^*)\lambda_k = 1_{M_2(\mathbb{C})}. \tag{8}$$

Hence, particularly for $k = 0, 1$, we have

$$\sum_j x_j x_j^* = \left(\begin{array}{cc} 1 & 0 \\ 0 & 1 \end{array}\right) \quad \text{and} \quad \sum_j x_j \alpha(x_j^*) = \sum_j x_j w x_j^* w^* = \left(\begin{array}{cc} 0 & 0 \\ 0 & 0 \end{array}\right).$$

With $x_j = \left(\begin{array}{cc} a_j & b_j \\ c_j & d_j \end{array}\right)$ and $z_i = e^{i\theta_i}$, $i = 1, 2$, this means

$$\sum_j \left(\begin{array}{cc} |a_j|^2 + |b_j|^2 & a_j\bar{c_j} + b_j\bar{d_j} \\ c_j\bar{a_j} + d_j\bar{b_j} & |c_j|^2 + |d_j|^2 \end{array}\right) = \left(\begin{array}{cc} 1 & 0 \\ 0 & 1 \end{array}\right),$$

$$\sum_j \left(\begin{array}{cc} |a_j|^2 + e^{i(\theta_2-\theta_1)}|b_j|^2 & e^{i(\theta_1-\theta_2)}a_j\bar{c_j} + b_j\bar{d_j} \\ |a_j|^2 + e^{i(\theta_2-\theta_1)}d_j\bar{b_j} & e^{i(\theta_1-\theta_2)}a_j\bar{c_j} + |d_j|^2 \end{array}\right) = \left(\begin{array}{cc} 0 & 0 \\ 0 & 0 \end{array}\right). \tag{9}$$

Therefore, by comparing (1,1) entries of each matrices, it follows that if α is saturated, then there exist positive real numbers $a\,(=\sum_j |a_j|^2) > 0$, $b\,(=\sum_j |b_j|^2) > 0$ such that

$$a + b = 1 \ \text{ and } a + e^{i(\theta_2-\theta_1)}b = 0. \tag{10}$$

Note that $b \neq 0$ since $b = 0$ implies $a = 0$ from $\sum_j \left(|a_j|^2 + e^{i(\theta_2-\theta_1)}d_j\bar{b_j}\right) = 0$ in (9). There are three possible cases for θ_1, θ_2 as follows.

(i) If $\theta_2 - \theta_1 \equiv 0 (\mathrm{mod}\ 2\pi)$, that is, α is trivial, then (10) is not possible.

(ii) If $\theta_2 - \theta_1 \equiv \pi (\mathrm{mod}\ 2\pi)$, then

$$x_1 = \frac{1}{\sqrt{2}}\left(\begin{array}{cc} 1 & 1 \\ 0 & 0 \end{array}\right) \quad \text{and} \quad x_2 = \frac{1}{\sqrt{2}}\left(\begin{array}{cc} 0 & 0 \\ 1 & 1 \end{array}\right)$$

satisfy (8) with $m = 2$. Thus α is saturated.

(iii) If $\theta_2 - \theta_1 \neq 0$, $\pi (\mathrm{mod}\ 2\pi)$, then (10) is not possible for any $a, b > 0$. Hence α is not saturated.

Remark 4.3. Let α, $\beta \in Aut(M)$ satisfy $\alpha^n = \beta^n = id_M$ for some $n \geq 1$. If there is a unitary $u \in M$ such that $\beta = Ad(u) \circ \alpha$, then α and β are said to be *exterior equivalent*, and if this is the case the crossed products are isomorphic, $M \times_\alpha G \cong M \times_\beta G$, [14, p.45]. Example 4.2 says that the property of being saturated may not be preserved under exterior equivalence. Also the case (iii) of Example 4.2 above with $w = diag(\lambda, \bar{\lambda})$, $\lambda = e^{\frac{2\pi}{3}}$ (hence $\theta_1 - \theta_2 = \frac{2\pi}{3} - \frac{4\pi}{3} \neq \pi (\mathrm{mod}\ 2\pi)$), shows that $Index(E) < |G|$ is possible even when E is of index-finite type. In fact, if $u_1 = \left(\begin{array}{cc} 1 & 0 \\ 0 & 1 \end{array}\right)$ and $u_2 = \left(\begin{array}{cc} 0 & 1 \\ 1 & 0 \end{array}\right)$, then $\{(u_i, u_i^*)\}_{i=1}^2$ forms a quasi-basis for E, but $Index(E) = 2 < |\mathbb{Z}_3|$.

Remark 4.4. Recall that the Rokhlin property and the tracial Rokhlin property (weaker than the Rokhlin property) are defined as follows and considered intensively in [7] and [12], respectively:

(a) ([7]) α is said to have the *Rokhlin property* if for every finite set $F \subset M$, every $\varepsilon > 0$, there are mutually orthogonal projections $\{e_g \mid g \in G\}$ in M such that
  (i) $\|\alpha_g(e_h) - e_{gh}\| < \varepsilon$ for $g, h \in G$.
  (ii) $\|e_g x - x e_g\| < \varepsilon$ for $g \in G$ and all $x \in F$.
  (iii) $\sum_{g\in G} e_g = 1$.

(b) ([12]) α is said to have the *tracial Rokhlin property* if for every finite set $F \subset M$, every $\varepsilon > 0$, every $n \in \mathbb{N}$, and every nonzero positive element $x \in M$, there are mutually orthogonal projections $\{e_g \mid g \in G\}$ in M such that:
   (i) $\|\alpha_g(e_h) - e_{gh}\| < \varepsilon$ for $g, h \in G$.
   (ii) $\|e_g x - x e_g\| < \varepsilon$ for $g \in G$ and all $x \in F$.
   (iii) With $e = \sum_{g\in G} e_g$, the projection $1 - e$ is Murray-von Neumann equivalent to a projection in the hereditary subalgebra of M generated by x.

The following proposition is actually observed in [12, Lemma 1.13], and we put a proof for reader's convenience.

Proposition 4.5. *Let M be a unital C^*-algebra and α be an action of a discrete group G on M. Suppose that for every $\varepsilon > 0$ and every finite subset $F \subset M$, there exist a family of projections $\{e_g\}_{g\in G}$ such that*
(1) $\|\alpha_g(e_h) - e_{gh}\| < \varepsilon$.
(2) $\|e_g x - x e_g\| < \varepsilon$ *for each* $x \in F$.
Then α is an outer action.

Proof. Suppose there is a unitary $u \in M$ such that $\alpha_g(x) = uxu^*$ for every $x \in M$ ($g \neq \iota$). Put $F = \{u\}$ and $0 < \varepsilon < 1/4$. Then there exist mutually orthogonal projections $\{e_g\}_{g\in G}$ such that $\|\alpha_g(e_h) - e_{gh}\| < \varepsilon < 1/4$. Thus $\|e_g u - u e_g\| < \varepsilon < 1/4$. Then $\|\alpha_g(e_\iota) - u e_\iota u^*\| = 0$. But

$$\begin{aligned} \|\alpha_g(e_\iota) - u e_\iota u^*\| &= \|\alpha_g(e_\iota) - e_g + e_g - e_\iota + e_\iota - u e_\iota u^*\| \\ &\geq \|e_g - e_\iota\| - \|\alpha_g(e_\iota) - e_g\| - \|e_\iota - u e_\iota 1 u^*\| \\ &\geq 1 - \frac{1}{4} - \frac{1}{4} = \frac{1}{2}, \end{aligned}$$

which is a contradiction. □

Remark 4.6. If $M \times_\alpha G$ is simple, α is obviously saturated, and this is the case if G is a finite group, M is α-simple, and $\tilde{\mathbb{T}}(\alpha_g) \neq \{1\}$ for all $g \neq \iota$ ([8, Theorem 3.1]). In particular, α is saturated if M is simple and α is outer.

But for a nonsimple M, this may not hold. In fact, if α is an outer action of $\mathbb{Z}_n$ on M and u is a unitary in M with $u^n = 1$ such that the action $Ad(u)$ on M is not saturated (as in Example 4.2), then the action $\alpha \oplus Ad(u)$ on $M \oplus M$ is outer but not saturated.

Now we show that if α satisfies the Rokhlin property (or satisfies the tracial Rokhlin property and M has cancellation) then α is saturated. For this we first review the cancellation property of C^*-algebras. For projections p, q in a C^*-algebra, we write $p \perp q$ if $pq = 0$, and $p \sim q$ if they are Murrey-von Neumann equivalent.

Definition 4.7. A unital C^*-algebra M has the *cancellation* if, whenever p, q, r are projections in $M_n(M)$ for some n, with $p \perp r$, $q \perp r$, and $(p + r) \sim (q + r)$, then $p \sim q$.

Remark 4.8. (1) If M has the cancellation and p, q are projections in M such that $(1-p) \sim (1-q)$, then $p \sim q$ ([4, V.2.4.14]).

(2) It is well known that every C^*-algebra with stable rank one has the cancellation ([4, V.3.1.24]).

Proposition 4.9. *Let α be an action of a finite group G on a unital C^*-algebra M. Then α is saturated if one of the following holds.*

(i) *α has the Rokhlin property.*

(ii) *α has the tracial Rokhlin property and M has the cancellation.*

Proof. (i) For an $\varepsilon > 0$, there exist mutually orthogonal projections $\{e_g\}_g$ such that $\sum_g e_g = 1$ and $\|\alpha_g(e_h) - e_{gh}\| < \varepsilon$. Then, with $m = 1$, the elements $b_g^1 := e_g$ satisfy (v) of Theorem 4.1.

(ii) Now suppose α has the tracial Rokhlin property and M has the cancellation. We shall show that $\mathcal{J}_\alpha$ contains the unit of $M \times_\alpha G$. Let $0 < \varepsilon < 1$. For each $g \in G$, choose mutually orthogonal projections $\{e_h^g\}_{h\in G}$ such that

$$\|\alpha_k(e_h^g) - e_{kh}^g\| < \frac{\varepsilon}{2|G|^2},$$

and put $e^g := \sum_{h\in G} e_h^g$. If $e^g = 1$, for some g, then $b_h^1 := e_h^g$ $(h \in G)$ will satisfy (v) of Theorem 4.1 as in (i). If $e^g \neq 1$ for every $g \in G$, then by the tracial Rokhlin property of α there exist mutually orthogonal projections $\{f_h^g\}_{h\in G}$ in M such that

$$\|\alpha_k(f_h^g) - f_{kh}^g\| < \frac{\varepsilon}{2|G|^2}$$

and

$$(1 - \sum_{h\in G} f_h^g) \sim (e^g)' < e^g$$

for a subprojection $(e^g)'$ of e^g. Put $f^g := \sum_h f_h^g$. Then since M has cancellation, it follows that $f^g \sim \big(1-(e^g)'\big) > (1-e^g)$. Let $v_g \in M$ be a partial isometry satisfying

$$v_g^* v_g = f^g, \quad v_g v_g^* = 1 - (e^g)',$$

and set

$$x_g := \frac{1}{|G|} \sum_{k,h} \Big((e_h^k \alpha_g(e_h^k) + (1-e^k) v_k f_h^k \alpha_g(f_h^k \alpha_{g^{-1}}(v_k^*)) \Big), \ g \in G.$$

Now we show that the element $x := \sum_g x_g \lambda_g \in \mathcal{J}_\alpha$ satisfies $\|x - 1\| < \varepsilon$. In fact, for $g \neq \iota$,

$$\begin{aligned}
\|x_g\| &\le \frac{1}{|G|} \sum_{k,h} \|(e_h^k \alpha_g(e_h^k) + (1-e^k) v_k f_h^k \alpha_g(f_h^k \alpha_{g^{-1}}(v_k^*))\| \\
&\le \frac{1}{|G|} \sum_{k,h} \big(\|(e_h^k \alpha_g(e_h^k)\| + \|f_h^k \alpha_g(f_h^k)\| \big) \\
&\le \frac{1}{|G|} \sum_{k,h} \big(\|e_h^k(\alpha_g(e_h^k) - e_{gh}^k)\| + \|e_h^k e_{gh}^k\| + \|f_h^k(\alpha_g(f_h^k) - f_{gh}^k)\| + \|f_h^k f_{gh}^k\| \big) \\
&\le \frac{1}{|G|} \sum_{k,h} (\frac{\varepsilon}{2|G|^2} + \frac{\varepsilon}{2|G|^2}) \\
&= \frac{\varepsilon}{|G|}
\end{aligned}$$

and

$$\begin{aligned} x_\iota &= \frac{1}{|G|}(\sum_k e^k + \sum_k (1-e^k)v_k f^k v_k^*) \\ &= \frac{1}{|G|}\big(\sum_k e^k + \sum_k (1-e^k)(1-(e^k)')\big) \\ &= 1. \end{aligned}$$

□

For the rest of this section we consider a finite group action on a commutative C^*-algebra $C(X)$. If G acts on a compact Hausdorff space X, it induces an action, say α, on $C(X)$ by

$$\alpha_g(f)(x) = f(g^{-1}x),\ f \in C(X).$$

For each $x \in X$, let $G_x = \{g \in G : gx = x\}$ be the isotropy group of x and for a subgroup H of G ($H < G$), put

$$X_H = \{x \in X : G_x = H\}.$$

It is readily seen that X_H and $X_{H'}$ are disjoint if $H \neq H'$, and X is partitioned as

$$X = \bigcup_{H<G} X_H.$$

Theorem 4.10. *Let X be a compact Hausdorff space and G a finite group acting on X. If α is the induced action of G on $C(X)$, the following are equivalent:*

(i) *$E : C(X) \to C(X)^\alpha$ is of index finite type.*
(ii) *X_H is closed for each $H < G$.*

Moreover, if this is case the index of E is $Index(E) = \sum_{H<G} \frac{|G|}{|H|}\chi_{X_H}$, where χ_{X_H} is the characteristic function on X_H.

Proof. (i) $\Longrightarrow$ (ii). If E is of index-finite type and $\{(u_i, u_i^*)\}_{i=1}^k$ is a quasi-basis for E, then

$$\sum_i u_i E(u_i^* f) = f,$$

that is,

$$\frac{1}{|G|}\sum_i u_i(x)\Big(\sum_{g\in G} u_i^*(g^{-1}x)f(g^{-1}x)\Big) = f(x), \tag{11}$$

for $f \in C(X)$ and $x \in X$. For each $x \in X$, choose a continuous function $f_x \in C(X)$ satisfying $f_x|_{Gx\setminus\{x\}} \equiv 0$ and $f_x(x) = 1$. Then (11) with f_x in place of f gives

$$\frac{1}{|G|}\sum_i u_i(x)\Big(\sum_{g\in G} u_i^*(g^{-1}x)f_x(g^{-1}x)\Big) = f_x(x), \tag{12}$$

and so we have

$$\frac{|G_x|}{|G|}\sum_i u_i(x)u_i^*(x) = 1. \tag{13}$$

To show that each X_H is closed, let $\{x_n \in X_H : n = 1, 2, \ldots\}$ be a sequence of elements in X_H with limit $x \in X_{H'}$. Then (11) gives

$$\begin{aligned} f_x(x_n) &= \frac{1}{|G|}\sum_i u_i(x_n)\Big(\sum_{g\in G} u_i^*(g^{-1}x_n)f_x(g^{-1}x_n)\Big) \\ &= \frac{1}{|G|}\sum_i u_i(x_n)\Big(\sum_{g\in H} u_i^*(g^{-1}x_n)f_x(g^{-1}x_n) + \sum_{g\notin H} u_i^*(g^{-1}x_n)f_x(g^{-1}x_n)\Big) \\ &= \frac{1}{|G|}\sum_i u_i(x_n)\Big(|H|u_i^*(x_n)f_x(x_n) + \sum_{g\notin H} u_i^*(g^{-1}x_n)f_x(g^{-1}x_n)\Big). \end{aligned}$$

Taking the limit as $n \to \infty$, we have

$$\begin{aligned} f_x(x) &= \frac{1}{|G|}\sum_i u_i(x)\Big(|H|u_i^*(x)f_x(x) + \sum_{g\notin H} u_i^*(g^{-1}x)f_x(g^{-1}x)\Big) \\ &= \frac{1}{|G|}\sum_i u_i(x)\Big(|H|u_i^*(x)f_x(x) + |H' \setminus H|u_i^*(x)f_x(x)\Big) \\ &= \frac{|H| + |H' \setminus H|}{|G|}\sum_i u_i(x)u_i^*(x)f_x(x). \end{aligned}$$

Therefore, comparing with (13), we obtain

$$|H'| = |H| + |H' \setminus H|$$

since $G_x = H'$ and $f_x(x) = 1$. Hence

$$H \subset H'.$$

On the other hand, since $G_{x_n} = H$, again by (13), $\frac{|H|}{|G|}\sum_i u_i(x_n)u_i^*(x_n) = 1$ with the limit $\frac{|H|}{|G|}\sum_i u_i(x)u_i^*(x) = 1$ as $n \to \infty$. But also $\frac{|H'|}{|G|}\sum_i u_i(x)u_i^*(x) = 1$ by (13), and thus $|H| = |H'|$. Consequently we have

$$H = H'$$

because $H \subset H'$. This shows that X_H is closed.

(ii) $\Longrightarrow$ (i). Assume that X_H is closed for every subgroup H of G. Then X_H is open since there are only finitely many such subsets. Let $\mathcal{U}_H = \{U_{H,i_H} : i_H = 1, 2, \ldots, n_H\}$ be an open covering of X_H such that

$$x \in U_{H,i_H} \implies g^{-1}x \notin U_{H,i_H} \text{ or } g^{-1}x \notin X_H \text{ whenever } g^{-1}x \neq x.$$

Let $\{v_{H,i_H}\}$ be a partition of unity subordinate to $\mathcal{U}_H$. We understand that the domain of v_{H,i_H} is X by assigning 0 to $x \notin X_H$. Let $u_{H,i_H} = \sqrt{v_{H,i_H}}$.

We claim that

$$\Big\{\Big(\sqrt{\frac{|G|}{|H|}}\,u_{H,i_H}, \sqrt{\frac{|G|}{|H|}}\,u_{H,i_H}^*\Big) : H < G, i_H = 1, 2, \ldots, n_H\Big\} \tag{14}$$

is a quasi-basis for E. For $f \in C(X)$ and $x \in X$, let $F < G$ and $1 \leq j \leq n_F$ be such that $x \in X_F$ and $x \in U_{F,j}$. Then

$$\begin{aligned}
&\sum_{H<G} \sum_{i_H=1}^{n_H} \Big(\sqrt{\frac{|G|}{|H|}}\, u_{H,i_H} E\big(\sqrt{\frac{|G|}{|H|}}\, u^*_{H,i_H} f \big) \Big)(x) \\
&= \frac{1}{|G|} \sum_{H<G} \sum_{i_H=1}^{n_H} \Big(\sqrt{\frac{|G|}{|H|}} u_{H,i_H}(x) \Big(\sum_{g \in G} \sqrt{\frac{|G|}{|H|}} u^*_{H,i_H}(g^{-1}x) f(g^{-1}x) \Big) \Big) \\
&= \sum_{i_F=1}^{n_F} \frac{1}{|F|} u_{F,i_F}(x) \Big(\sum_{g \in G} u^*_{F,i_F}(g^{-1}x) f(g^{-1}x) \Big) \\
&= \frac{1}{|F|} \sum_{i_F=1}^{n_F} u_{F,i_F}(x) \Big(\sum_{g \in F} u^*_{F,i_F}(g^{-1}x) f(g^{-1}x) \Big) \\
&= \frac{1}{|F|} \sum_{i_F=1}^{n_F} u_{F,i_F}(x) |F| u^*_{F,i_F}(x) f(x) \\
&= \sum_{i_F=1}^{n_F} v_{F,i_F}(x) f(x) \\
&= f(x),
\end{aligned}$$

as claimed.

□

Recall that an action G on X is *free* if $gx \neq x$ for $g \in G$, $g \neq 1$, and $x \in X$

Corollary 4.11. *Let X be a compact Hausdorff space and G a finite group acting on X. If α is the induced action on $C(X)$, the following are equivalent.*

- (i) *G acts freely on X.*
- (ii) *$E : C(X) \to C(X)^\alpha$ is of index-finite type with $Index(E) = |G|$.*
- (iii) *α is saturated.*

Proof. (i) $\Longrightarrow$ (ii) is proved in [19, Proposition 2.8.1].

To show (ii) $\Longrightarrow$ (i), let E be of index-finite type with $Index(E) = |G|$. Then from Theorem 4.10, we have

$$|G| = Index(E) = \sum_{H<G} \frac{|G|}{|H|} \chi_{X_H},$$

which implies that $H = \{\iota\}$ is the only subgroup of G such that $X_H \neq \emptyset$. Hence $X = X_{\{\iota\}}$, that is, G acts freely on X. (ii) $\Longleftrightarrow$ (iii) comes from Theorem 4.1. □

5. Saturated actions by compact groups

Notation 5.1. Let M be a C^*-algebra and α be an action of a compact group G on M. For $x, y \in M$, define continuous functions $f_{x,y}$, $f_{x,1}$, $f_{1,y} \in C(G, M)$ from G to

M as follows:

$$f_{x,y}(t) = x\alpha_t(y),$$
$$f_{x,1}(t) = x, \ \ f_{1,y}(t) = \alpha_t(y) \ \ \text{for } t \in G.$$

Then it is easily checked that $f_{x,y} = f_{x,1} * f_{1,y}$ and $f^*_{x,y} = f_{y^*,x^*}$.

Recall that $C(G, M)$ is a dense $*$-subalgebra of $M \times_\alpha G$ with the multiplication and involution defined by

$$f * g(t) = \int_G f(s)\alpha_s(g(s^{-1}t))ds,$$
$$f^*(t) = \alpha_t(f(t^{-1})^*),$$

where dg is the normalized Haar measure ([13, 7.7], [6, 8.3.1]). Hence if G is a finite group, $f_{x,y}$ can be written as $f_{x,y} = \frac{1}{|G|}\sum_g x\alpha_g(y)\lambda_g$.

If $\tilde{M}$ denotes the smallest unitization of M (so $\tilde{M} = M$ if M is unital), the function

$$e : G \to \tilde{M}, \ \ e(s) = 1, \ \ \text{for every } s \in G$$

is a projection of the multiplier algebra of $M \times_\alpha G$ ([17]).

Proposition 5.2. ([17]) *Let α be an action of a compact group G on a C^*-algebra M. Then identifying $x \in M^\alpha$ and the constant function in $C(G, M)$ with the value x everywhere we see that*

$$x \mapsto f_{x,1} : M^\alpha \to \ e(M \times_\alpha G)e$$

is an isomorphism of M^α onto the hereditary subalgebra $e(M \times_\alpha G)e$ of the crossed product $M \times_\alpha G$.

The notion of saturated action is introduced by Rieffel for a compact group action on a C^*-algebra, and we adopt the following equivalent condition as the definition.

Definition 5.3. (Rieffel, see [14, 7.1.9 Lemma]) Let M be a C^*-algebra and α be an action of compact group G on M. α is said to be *saturated* if the linear span of $\{f_{a,b} \mid a, b \in M\}$ is dense in $M \times_\alpha G$ (see Notation 5.1). We denote

$$\mathcal{J}_\alpha = \overline{span}\{f_{a,b} \mid a, b \in M\}.$$

Proposition 5.4. *Let α be an action of a compact group G on a C^*-algebra M. Then $\mathcal{J}_\alpha$ is the ideal of $M \times_\alpha G$ generated by the hereditary subalgebra $e(M \times_\alpha G)e$. Moreover $\mathcal{J}_\alpha = \overline{span}\{f_{a,a^*} \in C(G, M) \mid a \in M\}$.*

Proof. We first show that $\mathcal{J}_\alpha$ is an ideal of $M \times_\alpha G$. Let $x \in C(G, M)$ and $a, b \in M$. Then $x * f_{a,b} \in \mathcal{J}_\alpha$. Indeed,

$$\begin{aligned}(x * f_{a,b})(t) &= \int x(s)\alpha_s(f_{a,b}(s^{-1}t))ds \\ &= \int x(s)\alpha_s(a)\alpha_t(b)ds \\ &= \big(\int x(s)\alpha_s(a)ds\big)\alpha_t(b),\end{aligned}$$

hence $x * f_{a,b} = f_{c,b} \in \mathcal{J}_\alpha$, where $c = \int x(s)\alpha_s(a)ds \in M$. Also $f_{a,b}^* = f_{b^*,a^*}$ implies that $\mathcal{J}_\alpha = \mathcal{J}_\alpha{}^*$ is an ideal of $M \times_\alpha G$.

Let $\mathcal{J} := \overline{(M \times_\alpha G)e(M \times_\alpha G)}$ be the closed ideal generated by $e(M \times_\alpha G)e$. Now we show that $\mathcal{J}_\alpha \subset \mathcal{J}$. From

$$(f_{a,b} * e)(t) = \int f_{a,b}(s)\alpha_s(e(s^{-1}t))\, ds = \int a\alpha_s(b)\, ds = a \int \alpha_s(b)ds,$$

we have $f_{a,b} * e = f_{aE(b),1}$, where $E(b) = \int \alpha_s(b)ds \in M^\alpha$. Hence for a, b, c, and d in M, we have

$$\begin{aligned} f_{a,b} * e * f_{c,d} &= (f_{a,b} * e) * (f_{d^*,c^*} * e)^* \\ &= f_{aE(b),1} * (f_{d^*E(c^*),1})^* \\ &= f_{aE(b),1} * f_{1,E(c^*)d} \\ &= f_{aE(b),E(c^*)d}, \end{aligned}$$

which means that $f_{ax,yd} \in \mathcal{J}$ for any $a, d \in M$ and $x, y \in M^\alpha$. Since M^α contains an approximate identity for M, it follows that $f_{a,b} \in \mathcal{J}$ for $a, b \in A$.

For the converse inclusion $\mathcal{J} \subset \mathcal{J}_\alpha$, note that if $x \in C(G, M)$, then $(x * e)(t) = \int x(s)ds$ for $t \in G$. With notations $x' = \int x(s)ds$ and $x'' := \int \alpha_s(x(s^{-1}))ds\, (\in M)$, we see that

$$\begin{aligned} (x * e * y)(t) &= \int (x * e)(s)\alpha_s(y(s^{-1}t))ds \\ &= x' \int \alpha_s(y(s^{-1}t))ds \\ &= x'\alpha_t(\int \alpha_s(y(s^{-1}))ds) \\ &= x'\alpha_t(y'') \\ &= f_{x',y''}(t) \end{aligned}$$

belongs to $\mathcal{J}_\alpha$ for $x, y \in C(G, M)$.

Finally the following polarization identity proves the last assertion.

$$a\alpha_t(b) = \frac{1}{4}\sum_{k=0}^{3} i^k(b + i^k a^*)^*\alpha_t(b + i^k a^*).$$

□

6. THE GAUGE ACTION γ ON A GRAPH C^*-ALGEBRA

By a (directed) graph E we mean a quadruple $E = (E^0, E^1, r, s)$ consisting of the vertex set E^0, the edge set E^1, and the range, source maps $r, s : E^1 \to E^0$. If each vertex of E emits only finitely many edges E is called *row finite* and a row finite graph E is *locally finite* if each vertex receives only finitely many edges. By E^n we denote the set of all finite paths $\alpha = e_1 \cdots e_n$ $(r(e_i) = s(e_{i+1}), 1 \le i \le n-1)$ of *length* n $(|\alpha| = n)$. Each vertex is regarded as a finite path of length 0. Then $E^* = \cup_{n\ge 0} E^n$ is the set of all finite paths and the maps r and s naturally extend to E^*. A vertex v is called a *sink* if $s^{-1}(v) = \emptyset$ and a *source* if $r^{-1}(v) = \emptyset$.

If E is a row finite graph, we call a family $\{s_e, p_v \mid e \in E^1, v \in E^0\}$ of operators a *Cuntz-Krieger(CK) E-family* if $\{s_e\}_e$ are partial isometries and $\{p_v\}_v$ are mutually orthogonal projections such that

$$s_e^*s_e = p_{r(e)} \text{ and } p_v = \sum_{s(e)=v} s_e s_e^* \text{ if } s^{-1}(v) \neq \emptyset.$$

It is now well known that there exists a C^*-algebra $C^*(E)$ generated by a *universal* CK E-family $\{s_e, p_v \mid e \in E^1, v \in E^0\}$, in this case we simply write $C^*(E) = C^*(s_e, p_v)$. For the definition and basic properties of graph C^*-algebras, see, for example, [1, 2, 10, 11, 15] among others. If $\alpha = \alpha_1\alpha_2\cdots\alpha_{|\alpha|}$ ($\alpha_i \in E^1$) is a finite path, by s_α we denote the partial isometry $s_{\alpha_1}s_{\alpha_2}\cdots s_{\alpha_{|\alpha|}}$ ($s_v = s_v^* = p_v$, for $v \in E^0$).

We will consider only locally finite graphs and it is helpful to note the following properties of graph C^*-algebras.

Remark 6.1.
- (i) Let $C^*(E) = C^*(s_e, p_v)$ be the graph C^*-algebra associated with a row finite graph E, and let α, $\beta \in E^*$ be finite paths in E. Then

$$s_\alpha^* s_\beta = \begin{cases} s_\mu^*, & \text{if } \alpha = \beta\mu \\ s_\nu, & \text{if } \beta = \alpha\nu \\ 0, & \text{otherwise.} \end{cases}$$

  Therefore $C^*(E) = \overline{span}\{s_\alpha s_\beta^* \mid \alpha, \beta \in E^*\}$.
- (ii) Note that $s_\alpha s_\beta^* = 0$ for $\alpha, \beta \in E^*$ with $r(\alpha) \neq r(\beta)$.
- (iii) If α, β, μ, and ν in E^n are the paths of same length,

$$(s_\alpha s_\beta^*)(s_\mu s_\nu^*) = \delta_{\beta,\mu} s_\alpha s_\nu^*.$$

  Thus for each $n \in \mathbb{N}$ and a vertex v in a locally finite graph E, we see that

$$\overline{span}\{s_\alpha s_\beta^* \mid \alpha, \beta \in E^n \text{ and } r(\alpha) = r(\beta) = v\}$$

  is a $*$-algebra which is isomorphic to the full matrix algebra $M_m = (M_m(\mathbb{C}))$, where $m = \big|\{\alpha \in E^n \mid r(\alpha) = v\}\big|$.

Recall that the *gauge action* γ of $\mathbb{T}$ on $C^*(E) = C^*(s_e, p_v)$ is given by

$$\gamma_z(s_e) = zs_e, \ \gamma_z(p_v) = p_v, \quad z \in \mathbb{T}.$$

γ is well defined by the universal property of the CK E-family $\{s_e, p_v\}$. Since

$$\int_{\mathbb{T}} \gamma_z(s_\alpha s_\beta^*)dz = \int_{\mathbb{T}} z^{|\alpha|-|\beta|}(s_\alpha s_\beta^*)dz = 0, \ |\alpha| \neq |\beta|,$$

one sees that

$$C^*(E)^\gamma = \overline{span}\{s_\alpha s_\beta^* \mid \alpha, \beta \in E^*, \ |\alpha| = |\beta|\}.$$

If $\mathbf{Z}$ denotes the following graph:

$\mathbf{Z}$:



then $C^*(\mathbf{Z})$ is isomorphic to the C^*-algebra $\mathcal{K}$ of compact operators on an infinite dimensional separable Hilbert space, hence $C^*(\mathbf{Z})$ is itself a simple AF algebra. But $C^*(\mathbf{Z})^\gamma$ coincides with the commutative subalgebra $\overline{span}\{s_\alpha s_\alpha^* \mid \alpha \in \mathbf{Z}^*\}$ which is far from being simple, and thus we know that the simplicity of $C^*(E)$ does not imply that of $C^*(E)^\gamma$ in general.

In [9], the *Cartesian product* of two graphs E and F is defined to be the graph $E \times F = (E^0 \times F^0, E^1 \times F^1, r, s)$, where $r(e,f) = (r(e), r(f))$ and $s(e,f) = (s(e), s(f))$. Since the graph $\mathbf{Z} \times E$ has no loops for any row-finite graph E, we know that $C^*(E)^\gamma$ is an AF algebra ([10]) by the following proposition.

Proposition 6.2. ([9]) *Let E be a row finite graph with no sources. Then the following hold:*

(a) *$C^*(E)^\gamma$ is stably isomorphic to $C^*(E) \times_\gamma \mathbb{T}$.*
(b) *$C^*(E) \times_\gamma \mathbb{T} \cong C^*(\mathbf{Z} \times E)$.*

Now we show that a gauge action is saturated. For this, note that the linear span of the continuous functions of the form

$$t \mapsto f(t)x, \quad f \in C(G),\ x \in A$$

is dense in $C(G, A)$ [13, 7.6.1]. Hence by Remark 6.1(i), one sees that

$$C^*(E) \times_\gamma \mathbb{T} = \overline{span}\{z^n s_\alpha s_\beta^* \mid \alpha,\ \beta \in E^* \ \ n \in \mathbb{Z}\}. \tag{15}$$

Theorem 6.3. *Let E be a locally finite graph with no sinks and no sources. Then the gauge action γ on $C^*(E)$ is saturated.*

Proof. We show that $\mathcal{J}_\gamma = C^*(E) \times_\gamma \mathbb{T}$. By (15) it suffices to see that

$$z^n s_\alpha s_\beta^* \in \mathcal{J}_\gamma \quad \text{for all } \alpha, \beta \in E^*,\ n \geq 0$$

(because $z^{-n} s_\alpha s_\beta^* = (z^n s_\beta s_\alpha^*)^*$ for $n \geq 0$).

Now fix $\alpha,\ \beta \in E^*$ and $n \geq 0$. Put $l = n - (|\alpha| - |\beta|)$. There are two cases.

(i) $l \geq 0$: One can choose a path μ such that $|\mu| = l$ and $r(\mu) = s(\alpha)$. Then

$$z^n s_\alpha s_\beta^* = z^{l+|\alpha|-|\beta|} s_\mu^* s_\mu s_\alpha s_\beta^* = s_\mu^* \gamma_z(s_{\mu\alpha} s_\beta^*) = f_{s_\mu^*, s_{\mu\alpha} s_\beta^*}(z),$$

where the function $f_{s_\mu^*, s_{\mu\alpha} s_\beta^*}$ belongs to $\mathcal{J}_\gamma$.

(ii) $l < 0$: Choose a path ν with $|\nu| = |\beta| + n$ and $r(\nu) = r(\alpha)$. With $a = s_\alpha s_\nu^*$, $b = s_\nu s_\beta^*$, we have $f_{a,b} \in \mathcal{J}_\gamma$ and

$$z^n s_\alpha s_\beta^* = s_\alpha s_\nu^* \gamma_z(s_\nu s_\beta^*) = f_{a,b}(z).$$

□

Acknowledgements. The first author would like to thank Hiroyuki Osaka and Tamotsu Teruya for valuable discussions.

References

[1] T. Bates, J. H. Hong, I. Raeburn and W. Szymanski, *The ideal structure of the C^*-algebras of infinite graphs*, Illinois J. Math., **46**(2002), 1159–1176.

[2] T. Bates, D. Pask, I. Raeburn, and W. Szymanski, *The C^*-algebras of row-finite graphs*, New York J. Math. **6**(2000), 307–324.

[3] T. Bates and D. Pask, *Flow equivalence of graph algebras*, Ergod. Th. & Dynam. Sys. **24**(2004), 367–382.

[4] B. Blakadar, *Operator Algebras, Theory of C^*-algebras and von Neumann algebras*, Encyclopedia of mathematical sciences 122, Springer, 2006.

[5] J. Cuntz and W. Krieger, *A class of C^*-algebras and topological Markov chains*, Invent. Math. **56**(1980), 251–268.

[6] P. A. Fillmore, *A user's guide to operator algebras*, Canadian Math. Soc. Series of Monographs and Adv. Texts, John Wiley & Sons, Inc. 1996.

[7] M. Izumi, *Finite group actions on C^*-algebras with the Rohlin property, I*, Duke Math. J. **122**(2004), no. 2, 233–280.

[8] A. Kishimoto, *Outer automorphisms and related crossed products of simple C^*-algebras*, Comm. Math. Phys. **81**(1981), no. 1, 429–435.

[9] A. Kumjian and D. Pask, *C^*-algebras of directed graphs and group actions*, Ergodic Theory Dynam. Systems, **19**(1999), 1503–1519.

[10] A. Kumjian, D. Pask and I. Raeburn, *Cuntz-Krieger algebras of directed graphs*, Pacific J. Math. **184**(1998), 161–174.

[11] A. Kumjian, D. Pask, I. Raeburn and J. Renault *Graphs, groupoids, and Cuntz-Krieger algebras*, J. Funct. Anal. **144**(1997), 505–541.

[12] H. Osaka and N. C. Phillips, *Stable and real rank for crossed products by automorphisms with the tracial Rokhlin property*, Ergod. Th. & Dyn. Sys., to appear.

[13] G. K. Pedersen, *C^*-algebras and their automorphism groups*, London Math. Soc. Monographs No.14, Academic Press 1979.

[14] N. C. Phillips, *Equivariant K-theory and freness of group actions on C^*-algebras*, LNM 1274, Springer, 1987.

[15] I. Raeburn, Graph algebras, CBMS. 103, 2005, AMS.

[16] J. Renault, *A groupoid approach to C^*-algebras*, LNM 794, Springer, Berlin, 1980.

[17] J. Rosenberg, *Appendix to O. Bratteli's paper on "crossed products of UHF algebras*, Duke Math. J. **46**(1979), 25–26.

[18] W. Szymański and C. Peligrad, *Saturated actions of finite dimensional Hopf $*$-algebras on C^*-algebras*, Math. Scand. **75**(1994), 217–239.

[19] Y. Watatani, *Index for C^*-subalgebras*, Memoirs of the Amer. Math. Soc. 424, AMS, 1990.

[20] S. L. Woronowicz, *Compact matrix psedogroups*, Comm. Math. Phys. **111**(1987), 613–665.

Keywords: Finite dimensional Hopf $*$-algebra; saturated action; conditional expectation of index-finite type.

DEPARTMENT OF MATHEMATICAL SCIENCES AND RESEARCH INSTITUTE OF MATHEMATICS, SEOUL NATIONAL UNIVERSITY, SEOUL, 151–747, KOREA
E-mail address: `jajeong@snu.ac.kr`

DEPARTMENT OF MATHEMATICS, HANSHIN UNIVERSITY, OSAN, 447–791, KOREA
E-mail address: `ghpark@hanshin.ac.kr`